



A possible smeared phase transition in epidemic track-and-trace

Simon Dobson

School of Computer Science
University of St Andrews
Scotland UK

simon.dobson@st-andrews.ac.uk




University of
St Andrews | FOUNDED
1413 |



INTRODUCTION

Exploring epidemics using network science ¹

- ▶ Disease models and parameters
- ▶ Contact structures
- ▶ Countermeasures



In the course of writing about these ideas I stumbled across something I can't readily explain

- ▶ A change on behaviour that should be crisp, but doesn't seem to be
- ▶ Deserves more exploration

¹S. Dobson. *Epidemic modelling – Some notes, maths, and code*. Independent Publishing Network, 2020. ISBN 978-183853-565-0. URL <https://simoninireland.github.io/introduction-to-epidemics/>



STRUCTURE OF THIS TALK

Background

Measuring diseases

Compartmented models of disease

Epidemics on networks

Exploring diseases on networks

Changing the contact network

Immunity

Physical countermeasures


Track and trace

SEIR infections

Wild speculations



REAL DISEASES – 1




Different periods

- *Incubation*: from infection to onset of symptoms
- *Latent*: from exposure to infectiousness
- *Infectious*: overlapping with symptoms (usually)


Periods defined by biology, of both disease and host




REAL DISEASES – 2




Measles (Type A)



SARS (Type B)



Ebola (Type B)



Zombie apocalypse



\mathcal{R} AND ALL THAT ²

\mathcal{R} , the case reproduction number

- ▶ Number of secondary cases per primary
- ▶ Especially \mathcal{R}_0 , reproduction absent countermeasures

r , the case reproduction rate

- ▶ Doubling time for an epidemic
- ▶ Also sometimes see T_g , the inter-generation time

k , the overdispersion parameter

- ▶ Essentially a variance of the distribution around R
- ▶ Small number of individuals can drive most cases

²Royal Society SET-C group. Reproduction number (R) and growth rate (r) of the COVID-19 epidemic in the UK: methods of estimation, data sources, causes of heterogeneity, and use as a guide in policy formulation, August 2020. URL <https://royalsociety.org/-/media/policy/projects/set-c/set-covid-19-R-estimates.pdf>



THE “WICKEDNESS” OF COVID-19

$\mathcal{R}_0 \approx 3$ is not particularly infectious

- ▶ Straightforward to get $\mathcal{R} \approx 1.5$; harder to get $\mathcal{R} < 1$
- ▶ Significant overdispersion
- ▶ Infection seems to convey only temporary immunity

Case fatality rate is about 1%

- ▶ Too large to comfortably ignore, but not so large as to admit no arguments

Substantial asymptomatic transmission

- ▶ Asymmetric costs (spreading *vs* dying, “long covid”)
- ▶ Effective countermeasures are collective (and expensive)



THE GOALS OF MODELLING

What are we trying to find out?

- ▶ Concrete: how will this *particular* outbreak behave, in this *particular* population?
- ▶ Abstract: how can diseases behave *in general*? Are there common mathematical structures?



COMPARTMENTED MODELS

Traditional epidemic modelling uses the framework of a *compartmented model* of a disease

- ▶ A number of compartments holding some fraction of the population
- ▶ (Can also think of this as the state of each individual)
- ▶ Rules on how these fractions change over time



CONTINUOUS SIR

The model

- ▶ Susceptible individuals can catch the infection from *Infected* individuals
- ▶ ...who then are *Removed* from the dynamics by recovery (or death)


Epidemic dynamics

- ▶ Susceptibles infected per contact with probability β
- ▶ Infecteds removed with probability α

$$\frac{dS}{dt} = -\beta SI \qquad \frac{dI}{dt} = \beta SI - \alpha I \qquad \frac{dR}{dt} = \alpha I$$



SOLUTION





A RANGE OF MODELS WITH INCREASING COMPLEXITY

Can model different disease structures³

- ▶ SIR – simple infection offering complete immunity post-infection
- ▶ SIS – infection confers no immunity
- ▶ SEIR – exposed individuals are infectious before symptoms
- ▶ MSEIR – initial immunity passed from mother to child
- ▶ SEIRS – immunity wears off with time
- ▶ ...


³H. Hethcote. The mathematics of infectious diseases. *SIAM Review*, 42(4):599–653, December 2000. URL [doi://10.1137/S0036144500371907](https://doi.org/10.1137/S0036144500371907)



EPIDEMICS ON NETWORKS – 1

Complete homogeneous mixing implies populations have no structure, which clearly isn't true

- ▶ Not everyone meets everyone else
- ▶ Some are massively more (or less) connected than average





EPIDEMICS ON NETWORKS – 2

Use a network as the substrate for the epidemic ⁴

- ▶ Only nodes that are adjacent can interact
- ▶ Compartment = label on node

Leads to some more modelling decisions

- ▶ Degree distribution: probability p_k of node having degree k
- ▶ Often treat the mean degree $\langle k \rangle$ as “representative”
- ▶ Topological fine structure: loops, modules, layers, ...
- ▶ Adaptive behaviour to change these features in response to the progression of the disease (*e.g.*, quarantine)


⁴M. Newman. Spread of epidemic disease on networks. *Physical Review E*, 66, July 2002. URL [doi://10.1103/PhysRevE.66.016128](https://doi.org/10.1103/PhysRevE.66.016128)



HOW TO DO ANALYSIS

The “gold standard” is an analytic model with numerical validation

- ▶ Find an analytic description for what happens under different infection parameters
- ▶ Run process on random networks with the given topology
- ▶ Lots of repetitions to squeeze out variance
- ▶ (Hopefully) sample points land on solutions of the equations ⁵



⁵

P. Mann, V. A. Smith, J. Mitchell, and S. Dobson. Percolation in random graphs with higher-order clustering. URL <https://arxiv.org/abs/2006.06744>. Under review by Physical Review E. Preprint available on arXiv



HOW TO DO NUMERICAL VALIDATION

A networked population

- ▶ Set all to susceptible
- ▶ Change a fraction to infected, chosen at random
- ▶ Run the model rules until equilibrium

Discrete-event simulation ⁶

- ▶ After a time τ we see an event e (infection, removal, ...)
- ▶ Draw the next time and event from joint distribution $P(\tau, e)$

⁶D. Gillespie. Exact stochastic simulation of coupled chemical reactions. *Journal of Physical Chemistry*, 81(25): 2340—2361, 1977



FIELD LACKS STANDARD TOOLING

So we built some :-)

- ▶ epydemic: models, standard topologies, ...
- ▶ epyc: experiments, parallelism, dataset management

```
lab = epyc.ClusterLab(profile='hogun')

lab[epydemic.ERNetwork.N] = 10000
lab[epydemic.ERNetwork.KMEAN] = 40

lab[epydemic.SIR.P_INFECTED] = 0.001
lab[epydemic.SIR.P_REMOVE] = 0.002
lab[epydemic.SIR.P_INFECT] = numpy.linspace(0.00001, 0.0002, num=50)

m = epydemic.SIR()
g = epydemic.ERNetwork()
e = epydemic.StochasticDynamics(m, g)

lab.runExperiment(epyc.RepeatedExperiment(e, 100))
```



STRUCTURE OF THIS TALK

Background

- Measuring diseases

- Compartmented models of disease

- Epidemics on networks

Exploring diseases on networks

- Changing the contact network

- Immunity

- Physical countermeasures

Track and trace

- SEIR infections

- Wild speculations



WHAT DO HUMAN CONTACT NETWORKS LOOK LIKE?

Networks with particular contact structure

- ▶ Tend to have cycles (friends of friends)
- ▶ Highly variable numbers of contacts
- ▶ Often modular, having different local and global structures

Construction

- ▶ Theoretically, using a process to generate the structures
- ▶ Empirically, structured from survey data ⁷



⁷ K. Prem, A. Cook, and M. Jit. Projecting social contact matrices in 152 countries using contact surveys and demographic data. *PLOS Computational Biology*, 13(9), 2017. URL <https://doi.org/10.1371/journal.pcbi.1005697>



ER NETWORKS AND THE EPIDEMIC THRESHOLD

Erdős-Rényi (ER) networks

- For N nodes build the complete network K_N
- For each edge, retain (“occupy”) it with probability p_{infect}
- Leads to p_k normally distributed around $\langle k \rangle = p_{\text{infect}}N$







NOT ALL NETWORKS BEHAVE LIKE THIS

Too “even” to be a good model of human contacts

- Powerlaw with cutoff, $p_k \propto k^{-\alpha} e^{k/\kappa}$





- Relatively insensitive to p_{infect} , but sensitive to α and κ



HERD IMMUNITY

Sufficient immune/recovered individuals to stop an epidemic propagating

- ▶ Infecteds never adjacent to enough susceptibles
- ▶ First epidemic changes the effective topology
- ▶ “Effective” $\langle k \rangle$ drops from 20 to 5.5
- ▶ (Interested to explore other topological changes)





WHY THIS IS A BAD IDEA

Herd immunity has been seriously advocated as a strategy for COVID-19⁸

Ignores some rather inconvenient facts

- ▶ A 1% death rate = 700K UK deaths, about one year's excess
- ▶ At a rate that would collapse health services
- ▶ Immunity may not be permanent – which makes herd immunity behave differently (or not appear at all)
- ▶ Long COVID not accounted for in the costs

⁸See the “Great Barrington Declaration”, <https://gbdeclaration.org>



VACCINATION


“Herd immunity without the bad bits”

- ▶ Aim for the herd immunity threshold, generally about 60% of the population
- ▶ ...without anyone actually being infected

Epidemic proceeds at different rates depending on topology

- ▶ “Enough” contacts stabilise the size of outbreak

SIR over powerlaw networks for different cutoffs ($N = 10000$, $\alpha = 2$)





VACCINATION STRATEGIES


Randomly vaccinate 60% of the population

- ▶ Massive reduction in epidemic size
- ▶ Only catching high-degree nodes at random
- ▶ Sensitive to missing people


If we target vaccination we can reduce the threshold needed to get the same effect

- ▶ Target highest-degree 2% of nodes
- ▶ Take out the super-spreaders

SIR epidemic in the presence of 60% vaccination ($\kappa = 57.0$)



SIR epidemic vaccinating only the 2% highest-degree nodes ($N = 10000$, $\kappa = 57$)





PHYSICAL DISTANCING

What does a physically-distanced network look like?

- ▶ Good question
- ▶ (I have two SH projects looking at this at University level)

One possible model

- ▶ Normally-distributed, fully connected family “bubbles” of mean size 4
- ▶ A couple of members with outside contacts
- ▶ Exponentially-distributed connections between the contacts in different bubbles





LOCKDOWN CHANGES PROPAGATION

Changes the epidemic threshold compared to an ER network

- Needs a higher infectivity to take off

Slower take-off

- Not like a powerlaw network
- Get bursts of infection if the infection gets into a bubble





STRUCTURE OF THIS TALK

Background

- Measuring diseases

- Compartmented models of disease

- Epidemics on networks

Exploring diseases on networks

- Changing the contact network

- Immunity

- Physical countermeasures

Track and trace

- SEIR infections

- Wild speculations



ASYMPTOMATIC TRANSMISSION

Because COVID-19 is essentially SEIR (or maybe SEIRS) it invites other countermeasures

- ▶ Self-isolating on showing symptoms is ineffective
- ▶ Try to find the asymptomatic carriers

This is the basis for track-and-trace

- ▶ Identify contacts of that person
- ▶ Quarantine them if they're infected – means we catch infecting individuals before they knew to self-isolate
- ▶ Quarantine the symptomatic individual too



TRACK AND TRACE IN PRACTICE

A large-scale procedure, unlike the local procedure of self-isolation when symptomatic

- ▶ Requires organisation by some authority
- ▶ What can possibly go wrong?...

Unlikely to be fully accurate even if done competently


- ▶ Some proportion of people don't quarantine? (p_{rewire})
- ▶ Only test some proportion of contacts? (p_{detect})



THE IMPACT OF DETECTION RATES

Hold p_{rewire} constant and vary p_{detect}

SEIR epidemic size vs p_{detect} ($N = 10000$, $\langle k \rangle = 40$, $p_{rewire} = 0.8$)





INTERPRETING THE PLOT


Very dependent on detection rate

- ▶ Quarantine can still sometimes damp epidemic anyway
- ▶ High detection is very effective

Seems to be unstable in the mid range

- ▶ All sizes of epidemic are possible
- ▶ No clean break
- ▶ Phase transition is “smeared out” rather than occurring cleanly

SEIR epidemic size vs p_{detect} ($N = 10000$, $\langle k \rangle = 40$, $p_{\text{rewire}} = 0.8$)





WHAT'S HAPPENING?

This phenomenon only entered the literature last year ⁹

Possible explanations

- ▶ It's an artefact of ER networks
- ▶ It's an artefact of not simulating long enough to crush the variance
- ▶ It's a function of network fine structure like clustering
- ▶ SEIR propagation is really driven by asymptomatic infection

⁹L. Hébert-Dufresne and A. Allard. Smeared phase transitions in percolation on real complex networks. *Physical Review Research*, 1, August 2019. URL <https://doi.org/10.1103/PhysRevResearch.1.013009>



POSSIBLE IMPLICATIONS

Designing track-and-trace

- ▶ Very sensitive to the proportion of contacts tested
- ▶ Need to check at least 40% to have any effect at all

Effectiveness

- ▶ Even very effective tracing doesn't guarantee elimination
- ▶ Although it does reduce the peak significantly once above about 80%
- ▶ Understanding the smearing might let us improve the test strategy



FOUR THINGS TO TAKE AWAY

1. Epidemic spreading still isn't fully understood, even now – there's lots of exciting work still to do, mathematically and computationally
2. Explore of possible public policy decisions
3. Need (in my copious free time) to understand what's going on with this possible smeared phase transition
4. Especially interested in how small-scale topological structures affect network-based processes



REFERENCES



S. Dobson. *Epidemic modelling – Some notes, maths, and code*. Independent Publishing Network, 2020. ISBN 978-183853-565-0. URL <https://simoninireland.github.io/introduction-to-epidemics/>.



D. Gillespie. Exact stochastic simulation of coupled chemical reactions. *Journal of Physical Chemistry*, 81(25): 2340—2361, 1977.



L. Hébert-Dufresne and A. Allard. Smeared phase transitions in percolation on real complex networks. *Physical Review Research*, 1, August 2019. URL <https://doi.org/10.1103/PhysRevResearch.1.013009>.



H. Hethcote. The mathematics of infectious diseases. *SIAM Review*, 42(4):599–653, December 2000. URL [doi://10.1137/S0036144500371907](https://doi.org/10.1137/S0036144500371907).



P. Mann, V. A. Smith, J. Mitchell, and S. Dobson. Percolation in random graphs with higher-order clustering. URL <https://arxiv.org/abs/2006.06744>. Under review by Physical Review E. Preprint available on arXiv.



M. Newman. Spread of epidemic disease on networks. *Physical Review E*, 66, July 2002. URL [doi://10.1103/PhysRevE.66.016128](https://doi.org/10.1103/PhysRevE.66.016128).



K. Prem, A. Cook, and M. Jit. Projecting social contact matrices in 152 countries using contact surveys and demographic data. *PLOS Computational Biology*, 13(9), 2017. URL <https://doi.org/10.1371/journal.pcbi.1005697>.



Royal Society SET-C group. Reproduction number (R) and growth rate (r) of the COVID-19 epidemic in the UK: methods of estimation, data sources, causes of heterogeneity, and use as a guide in policy formulation, August 2020. URL <https://royalsociety.org/-/media/policy/projects/set-c/set-covid-19-R-estimates.pdf>.